arXiv:1109.3930v1 [math.CO] 19 Sep 2011

Roman Bondage Number of a Graph[*]

Fu-Tao Hu, Jun-Ming Xu[†]

Department of Mathematics
University of Science and Technology of China
Hefei, Anhui, 230026, China

Abstract: The Roman dominating function on a graph $G = (V, E)$ is a function $f : V \to \{0, 1, 2\}$ such that each vertex x with $f(x) = 0$ is adjacent to at least one vertex y with $f(y) = 2$. The value $f(G) = \sum\limits_{u \in V(G)} f(u)$ is called the weight of f. The Roman domination number $\gamma_{\rm R}(G)$ is defined as the minimum weight of all Roman dominating functions. This paper defines the Roman bondage number $b_{\rm R}(G)$ of a nonempty graph $G = (V, E)$ to be the cardinality among all sets of edges $B \subseteq E$ for which $\gamma_{\rm R}(G - B) > \gamma_{\rm R}(G)$. Some bounds are obtained for $b_{\rm R}(G)$, and the exact values are determined for several classes of graphs. Moreover, the decision problem for $b_{\rm R}(G)$ is proved to be NP-hard even for bipartite graphs.

Keywords: Roman domination number, Roman bondage number,NP-hardness.

AMS Subject Classification: 05C69

1 Introduction

In this paper, a graph $G = (V, E)$ is considered as an undirected graph without loops and multi-edges, where $V = V(G)$ is the vertex set and $E = E(G)$ is the edge set. For each vertex $x \in V(G)$, let $N_G(x) = \{y \in V(G) : (x, y) \in E(G)\}$, $N_G[x] = N_G(x) \cup \{x\}$.

---

[*] The work was supported by NNSF of China (No. 11071233).

[†] Corresponding author: xujm@ustc.edu.cn

A subset $S \subseteq V$ is a *dominating set* of G if $N_G[x] \cap S \neq \emptyset$ for every vertex x in G. The *domination number* of G, denoted by $\gamma(G)$, is the minimum cardinality of all dominating sets of G. The *Roman dominating function* on a graph $G = (V, E)$, proposed by Cockayne *et al.* [2], is a function $f : V \to \{0, 1, 2\}$ such that each vertex x with $f(x) = 0$ is adjacent to at least one vertex y with $f(y) = 2$. Let (V_0, V_1, V_2) be the ordered partition of V induced by f, where $V_i = \{v \in V | f(v) = i\}$ for $i = 0, 1, 2$. It is clear that $V_1 \cup V_2$ is a dominating set of G, called *the Roman dominating set*, denoted by $D_{\rm R} = (V_1, V_2)$. For $S \subseteq V$, let $f(S) = \sum\limits_{u \in S} f(u)$. The value $f(V(G))$ is called the *weight* of f, denoted by $f(G)$. The *Roman domination number*, denoted by $\gamma_{\rm R}(G)$, is defined as the minimum weight of all Roman dominating functions, that is,

$$\gamma_{\rm R}(G) = \min\{f(G) : f \text{ is a Roman dominating function on } G\}.$$

It is clear that for a Roman dominating function f on G and a Roman dominating set $D_{\rm R}$ of G, $f(D_{\rm R}) = 2|V_2| + |V_1|$. If $D_{\rm R}$ is a minimum Roman dominating set of graph G, then $f(D_{\rm R}) = \gamma_{\rm R}(G)$. A Roman dominating function f is called a $\gamma_{\rm R}$-*function* if $f(G) = \gamma_{\rm R}(G)$. It has been showed by Cockayne *et al.* [2] that for any graph G, $\gamma(G) \leqslant \gamma_R(G) \leqslant 2\gamma(G)$. A graph G is called to be *Roman* if $\gamma_{\rm R}(G) = 2\gamma(G)$. Roman domination numbers have been studied, for example, in [2, 3, 5, 6, 11, 12, 13, 14, 15, 16, 20].

To measure the vulnerability or the stability of the domination in an interconnection network under edge failure, Fink et at. [4] proposed the concept of the bondage number in 1990. The *bondage number*, denoted by $b(G)$, of G is the minimum number of edges whose removal from G results in a graph with larger domination number of G.

Analogously, we can define the Roman bondage number. The *Roman bondage number*, denoted by $b_{\rm R}(G)$, of a nonempty graph G is the minimum number of edges whose removal from G results in a graph with larger Roman domination number. Precisely speaking, the Roman bondage number

$$b_{\rm R}(G) = \min\{|B| : B \subseteq E(G), \gamma_{\rm R}(G - B) > \gamma_{\rm R}(G)\}.$$

An edge set B that $\gamma_{\rm R}(G - B) > \gamma_{\rm R}(G)$ is called the *Roman bondage set* and the minimum one the *minimum Roman bondage set.* In fact, if B is a minimum Roman bondage set, then $\gamma_{\rm R}(G-B)=\gamma_{\rm R}(G)+1$, because the removal of one single edge can not increase the Roman domination number by more than one. If $b_{\rm R}(G)$ does not exist we define $b_{\rm R}(G)=\infty$.

In this paper, we give an original investigation. Some bounds are obtained for $b_{\rm R}(G)$, and the exact values are determined for several classes of graphs. Moreover, the decision problem for $b_{\rm R}(G)$ is proved to be NP-hard even for bipartite graphs.

In the proofs of our results, when a Roman dominating function of a graph is constructed, we only give its nonzero value of some vertices.

2 Some basic results on $\gamma_{\rm R}$

For terminology and notation on graph theory not given here, the reader is referred to [8, 9, 19].

Let $G=(V,E)$ be a graph and $E_G(x)=\{xy\in E(G): y\in N_G(x)\}$. For two disjoint nonempty sets $S,T\subset V(G)$, $E_G(S,T)=E(S,T)$ denotes the set of edges between S and T. The degree of x is denoted by $d_G(x)$, which is equal to $|N_G(x)|$, and n_i denotes the number of vertices of degree i in G for $i=1,2,\cdots,\Delta(G)$. Denote the maximum and the minimum degree of G by $\Delta(G)$ and $\delta(G)$, respectively.

The symbols P_n and C_n denote a path and a cycle, respectively, where $V(P_n)=V(C_n)=\{x_1,x_2,\cdots,x_n\}$, $E(P_n)=\{x_ix_{i+1}:\ i=1,2,\cdots,n\}$ and $E(C_n)=E(P_n)\cup\{x_1x_n\}$.

The Cartesian product graph $G_1\times G_2$ of two graphs $G_1=(V_1,E_1)$ and $G_2=(V_2,E_2)$ is a graph with vertex-set $V=V_1\times V_2=\{(x,y):\ x\in V_1, y\in V_2\}$, and two vertices (x_1,y_1) and (x_2,y_2) being adjacent if and only if either $x_1=x_2$, y_1 and y_2 are adjacent in G_2, or $y_1=y_2$, x_1 and x_2 are adjacent in G_1.

In this section, we recall some basic results on $\gamma_{\rm R}$, which will be used in our discussion.

Lemma 2.1 (Cockayne et al. [2]) *For a path P_n and a cycle C_n,*

$$\gamma_{\rm R}(P_n)=\gamma_{\rm R}(C_n)=\left\lceil\frac{2n}{3}\right\rceil.$$

For a grid graph $P_2 \times P_n$*,*

$$\gamma_{\rm R}(P_2 \times P_n) = n + 1.$$

For a complete t*-partite graph* $K_{m_1,m_2,\cdots,m_t}$ *with* $1 \leq m_1 \leq m_2 \leq \cdots \leq m_t$ *and* $t \geq 2$*,*

$$\gamma_{\rm R}(K_{m_1,m_2,\cdots,m_t}) = \begin{cases} 2, & \text{if } m_1 = 1; \\ 3, & \text{if } m_1 = 2; \\ 4, & \text{if } m_1 \geq 3. \end{cases}$$

Lemma 2.2 (Cockayne et al. [2]) *If* G *is a graph of order* n *and contains vertices of degree* $n-1$*, then* $\gamma_{\rm R}(G) = 2$*.*

Lemma 2.3 *Let* G *be a nonempty graph with order* $n \geq 3$*, then* $\gamma_{\rm R}(G) = 3$ *if and only if* $\Delta(G) = n - 2$*.*

Proof. Assume that u is a vertex of degree $n-2$ and v is the unique vertex not adjacent to u in G. It is easy to verify that $\gamma_{\rm R}(G) \geq 3$. Let f be a function from $V(G)$ to $\{0,1,2\}$ subject to

$$f(x) = \begin{cases} 2, & \text{if } x = u; \\ 1, & \text{if } x = v; \\ 0, & \text{otherwise.} \end{cases}$$

Then f is a Roman dominating function of G with $f(G) = 3$. Thus, $\gamma_{\rm R}(G) = 3$.

Conversely, assume $\gamma_{\rm R}(G) = 3$, then $\Delta(G) \leq n - 2$ by Lemma 2.2. Let f be a $\gamma_{\rm R}$-function of G.

If there is no vertex u with $f(u) = 2$, then $f(v) = 1$ for each vertex $v \in V(G)$, and so $n = 3$ since $f(G) = \gamma_{\rm R}(G) = 3$. Sine G is nonempty and not K_3, G consists of K_2 and an isolated vertex. Thus, $\Delta(G) = 1 = n - 2$.

If there is a vertex u with $f(u) = 2$, then there is only one vertex $v \in V(G)$ with $f(v) = 1$ since $f(G) = \gamma_{\rm R}(G) = 3$. The other $n-2$ vertices assigned 0 are all adjacent to u. Thus, $\Delta(G) \geq d_G(u) \geq n-2$ and hence $\Delta(G) = n-2$. ∎

Lemma 2.4 *Let* G *be an* $(n-3)$*-regular graph with order* $n \geq 4$*. Then* $\gamma_{\rm R}(G) = 4$*.*

Proof. Since G is $(n-3)$-regular and $n \geq 4$, G is nonempty. It is clear that $\gamma_{\rm R}(G) > 2$. By Lemma 2.3, $\gamma_{\rm R}(G) \neq 3$ since $\Delta(G) = n - 3$. Then $\gamma_{\rm R}(G) \geq 4$. For any vertex

$x \in V(G)$, let y, z be the only two vertices not adjacent to x in G, let $f(x) = 2$ and $f(y) = f(z) = 1$. Then, f is a Roman dominating function of G with $f(G) = 4$, hence $\gamma_{\mathrm{R}}(G) \leq 4$. Thus, $\gamma_{\mathrm{R}}(G) = 4$. ∎

Lemma 2.5 (Cockayne et al. [2]) *For any graph G, $\gamma(G) \leq \gamma_{\mathrm{R}}(G) \leq 2\gamma(G)$.*

Lemma 2.6 (Cockayne et al. [2]) *For any graph G of order n, $\gamma(G) = \gamma_{\mathrm{R}}(G)$ if and only if $G = \bar{K}_n$.*

Lemma 2.7 (Cockayne et al. [2]) *If G is a connected graph of order n, then $\gamma_{\mathrm{R}}(G) = \gamma(G) + 1$ if and only if there is a vertex $v \in V(G)$ of degree $n - \gamma(G)$.*

A graph G is called to be *vertex domination-critical* (*vc-graph* for short) if $\gamma(G-x) < \gamma(G)$ for any vertex x in G.

Lemma 2.8 (Brigham, Chinn and Dutton [1], 1988) *A graph G with $\gamma(G) = 2$ is a vc-graph if and only if G is a complete graph $K_{2t}\,(t \geq 2)$ with a perfect matching removed.*

A graph G of order n is *vertex Roman domination-critical* (*vrc-graph* for short) if $\gamma_{\mathrm{R}}(G) \neq n$ and $\gamma_{\mathrm{R}}(G - x) < \gamma_{\mathrm{R}}(G)$ for any vertex x in G. For example, for a positive integer k, both C_{3k+1} and C_{3k+2} are vrc-graphs by Lemma 2.1. From the definition, it is clear that $\gamma_{\mathrm{R}}(G) \geq 3$ if G is a vrc-graph with order at least 3.

Lemma 2.9 *If G is a vrc-graph with $\gamma_{\mathrm{R}}(G) = 3$, then G is a vc-graph with $\gamma(G) = 2$.*

Proof. Let G be a vrc-graph with $\gamma_{\mathrm{R}}(G) = 3$. From the definition of vrc-graph, $|V(G)| > \gamma_{\mathrm{R}}(G) = 3$. By Lemma 2.3, $\Delta(G) = |V(G)| - 2$ and hence $\gamma(G) = 2$. For any vertex x, if $\gamma_{\mathrm{R}}(G - x) < \gamma_{\mathrm{R}}(G) = 3$, then $\gamma_{\mathrm{R}}(G - x) = 2$ since $G - x$ is nonempty. By Lemma 2.2, $G - x$ contains vertices of degree $|V(G - x)| - 1$ and, hence, $\gamma(G - x) = 1$, which implies that G is a vc-graph. ∎

3 The exact values of $b_{\rm R}$ for some graphs

Lemma 3.1 *Let G be a graph with order $n \geq 3$ and t be the number of vertices of degree $n-1$ in G. If $t \geq 1$ then $b_{\rm R}(G) = \lceil \frac{t}{2} \rceil$.*

Proof. Let H be a spanning subgraph of G obtained by removing fewer than $\lceil \frac{t}{2} \rceil$ edges from G. Then H contains vertices of degree $n-1$ and, hence, $\gamma_{\rm R}(H) = 2 = \gamma_{\rm R}(G)$ by Lemma 2.2, which implies $b_{\rm R}(G) \geq \lceil \frac{t}{2} \rceil$.

Since G contains t vertices of degree $n-1$, it contains a complete subgraph K_t induced by these t vertices. We can remove $\lceil \frac{t}{2} \rceil$ edges such that no vertices have degree $n-1$ and, hence, $\gamma_{\rm R}(H) \geq 3 > 2 = \gamma_{\rm R}(G)$ since $n \geq 3$. Thus $b_{\rm R}(G) \leq \lceil \frac{t}{2} \rceil$, whence $b_{\rm R}(G) = \lceil \frac{t}{2} \rceil$. ∎

Corollary 3.1 *For a complete graph K_n $(n \geq 3)$, $b_{\rm R}(K_n) = \lceil \frac{n}{2} \rceil$.*

Theorem 3.1 *For a path P_n with $n \geq 3$,*

$$b_{\rm R}(P_n) = \begin{cases} 1, & \text{if } n \equiv 0, 1 \pmod 3; \\ 2, & \text{otherwise.} \end{cases}$$

Proof. Let $P_n = (x_1, x_2, \ldots, x_n)$ be a path. By Lemma 2.1, $\gamma_{\rm R}(P_n) = \lceil \frac{2n}{3} \rceil$.

If $n \equiv 0, 1 \pmod 3$, then

$$\gamma_{\rm R}(P_n - x_2x_3) = 2 + \left\lceil \frac{2(n-2)}{3} \right\rceil = 1 + \left\lceil \frac{2n-1}{3} \right\rceil = 1 + \gamma_{\rm R}(P_n),$$

and hence $b_{\rm R}(P_n) \leq 1$, whence $b_{\rm R}(P_n) = 1$.

If $n \equiv 2 \pmod 3$, then for any edge $e = x_ix_{i+1} \in E(P_n)$,

$$\gamma_{\rm R}(P_n - e) = \left\lceil \frac{2i}{3} \right\rceil + \left\lceil \frac{2(n-i)}{3} \right\rceil \leq \left\lceil \frac{2(n-i)+2i+2}{3} \right\rceil = \left\lceil \frac{2n}{3} \right\rceil = \gamma_{\rm R}(P_n),$$

and hence $b_{\rm R}(P_n) \geq 2$. Since

$$\gamma_{\rm R}(P_n - x_2x_3 - x_4x_5) = 2 + 2 + \left\lceil \frac{2(n-4)}{3} \right\rceil = 1 + \left\lceil \frac{2n+1}{3} \right\rceil \geq 1 + \gamma_{\rm R}(P_n),$$

we have $b_{\rm R}(P_n) \leq 2$, whence $b_{\rm R}(P_n) = 2$. ∎

Corollary 3.2 *For a cycle C_n with $n \geq 3$,*

$$b_{\rm R}(C_n) = \begin{cases} 2, & \text{if } n = 0, 1 \pmod 3; \\ 3, & \text{otherwise.} \end{cases}$$

Lemma 3.2 *Let $P_n = (x_1, x_2, \ldots, x_n)$ be a path, and use $u_{i,j}$ to denote the vertex (x_i, x_j) in $P_2 \times P_n$, where $1 \leq i \leq 2$ and $1 \leq j \leq n$. Then there exists a $\gamma_{\rm R}$-function f on $P_2 \times P_n$ such that $f(u_{1,1}) = 2$ or $f(u_{2,1}) = 2$ or $f(u_{1,n}) = 2$ or $f(u_{2,n}) = 2$.*

Proof. Without loss of generality, we only need to find a $\gamma_{\rm R}$-function f on $P_2 \times P_n$ with $f(u_{1,1}) = 2$. Define a Roman dominating function f as follows. For each non-negative integer i with $1 + 4i \leq n$, let $f(u_{1,1+4i}) = 2$, and for each non-negative integer j with $3 + 4j \leq n$, let $f(u_{2,3+4j}) = 2$. If $n \equiv 0 \pmod 4$, let $f(u_{1,n}) = 1$, and if $n \equiv 2 \pmod 4$, let $f(u_{2,n}) = 1$. Then $f(P_2 \times P_n) = n + 1$ and, hence by Lemma 2.1, f is a $\gamma_{\rm R}$-function with $f(u_{1,1}) = 2$. ∎

Theorem 3.2 *$b_{\rm R}(P_2 \times P_n) = 2$ for $n \geq 2$.*

Proof. By Lemma 2.1, we have $\gamma_{\rm R}(P_2 \times P_n) = n + 1$. Since $\gamma_{\rm R}(P_2 \times P_n - u_{1,1}u_{1,2} - u_{2,1}u_{2,2}) = 2 + \gamma_{\rm R}(P_2 \times P_{n-1}) = n + 2$, we have $b_{\rm R}(P_2 \times P_n) \leq 2$. Next we prove that $\gamma_{\rm R}(P_2 \times P_n - e) \leq \gamma_{\rm R}(P_2 \times P_n)$ for any edge $e \in E(P_2 \times P_n)$.

Suppose that e is incident with some vertex in $\{u_{1,1}, u_{2,1}, u_{1,n}, u_{2,n}\}$. Without loss of generality let e be incident with $u_{1,1}$. By Lemma 3.2, there exists a $\gamma_{\rm R}$-function f on $P_2 \times (P_n - P_1)$ such that $f(u_{2,2}) = 2$. Denote $f(u_{1,1}) = 1$ and then f is a Roman dominating function of $P_2 \times P_n - e$ with $f(P_2 \times P_n - e) = n + 1$, thus $\gamma_{\rm R}(P_2 \times P_n - e) \leq \gamma_{\rm R}(P_2 \times P_n)$.

Suppose that e is incident with some vertex in $\{u_{i,j} : 1 \leq i \leq 2, 2 \leq j \leq n - 1\} \setminus \{u_{1,1}, u_{2,1}, u_{1,n}, u_{2,n}\}$. Without loss of generality let e be incident with $u_{1,j}$ and not incident with $u_{1,j-1}$. By Lemma 3.2, there exists a $\gamma_{\rm R}$-function f_1 on $P_2 \times P_{j-1}$ with $f_1(u_{1,j-1}) = 2$ and a $\gamma_{\rm R}$-function f_2 on $P_2 \times (P_n - P_j)$ with $f_2(u_{2,j+1}) = 2$. Then $f = f_1 \cup f_2$ is a Roman dominating function on $P_2 \times P_n - e$ with $f(P_2 \times P_n - e) = n + 1$, thus $\gamma_{\rm R}(P_2 \times P_n - e) \leq \gamma_{\rm R}(P_2 \times P_n)$.

The above two cases yield that $b_{\rm R}(P_2 \times P_n) \geq 2$ and, hence, $b_{\rm R}(P_2 \times P_n) = 2$. The lemma follows. ∎

4 Complexity of Roman bondage number

In this section, we will show that the Roman bondage number problem is NP-hard and the Roman domination number problem is NP-complete even for bipartite graphs. We first state the problem as the following decision problem.

Roman bondage number problem (RBN):

Instance: *A nonempty bipartite graph G and a positive integer k.*

Question: *Is $b_{\rm R}(G) \le k$?*

Roman domination number problem (RDN):

Instance: *A nonempty bipartite graph G and a positive integer k.*

Question: *Is $\gamma_{\rm R}(G) \le k$?*

Following Garey and Johnson's techniques for proving NP-completeness given in [7], we prove our results by describing a polynomial transformation from the known-well NP-complete problem: 3SAT. To state 3SAT, we recall some terms.

Let U be a set of Boolean variables. A *truth assignment* for U is a mapping $t: U \to \{T, F\}$. If $t(u) = T$, then u is said to be "true" under t; If $t(u) = F$, then u is said to be "false" under t. If u is a variable in U, then u and $\bar{u}$ are *literals* over U. The literal u is true under t if and only if the variable u is true under t; the literal $\bar{u}$ is true if and only if the variable u is false.

A *clause* over U is a set of literals over U. It represents the disjunction of these literals and is *satisfied* by a truth assignment if and only if at least one of its members is true under that assignment. A collection $\mathscr{C}$ of clauses over U is *satisfiable* if and only if there exists some truth assignment for U that simultaneously satisfies all the clauses in $\mathscr{C}$. Such a truth assignment is called a *satisfying truth assignment* for $\mathscr{C}$. The 3SAT is specified as follows.

3-satisfiability problem (3SAT):

Instance: *A collection $\mathscr{C} = \{C_1, C_2, \ldots, C_m\}$ of clauses over a finite set U of variables such that $|C_j| = 3$ for $j = 1, 2, \ldots, m$.*

Question: *Is there a truth assignment for U that satisfies all the clauses in $\mathscr{C}$?*

Theorem 4.1 (Theorem 3.1 in [7]) *3SAT is NP-complete.*

Theorem 4.2 *RBN is NP-hard even for bipartite graphs.*

Proof. The transformation is from 3SAT. Let $U = \{u_1, u_2, \ldots, u_n\}$ and $\mathscr{C} = \{C_1, C_2, \ldots, C_m\}$ be an arbitrary instance of 3SAT. We will construct a bipartite graph G and choose an integer k such that $\mathscr{C}$ is satisfiable if and only if $b_{\rm R}(G) \le k$. We construct such a graph G as follows.

For each $i = 1, 2, \ldots, n$, corresponding to the variable $u_i \in U$, associate a graph H_i with vertex set $V(H_i) = \{u_i, \bar{u}_i, v_i, v'_i, x_i, y_i, z_i, w_i\}$ and edge set $E(H_i) = \{u_iv_i, u_iz_i, \bar{u}_iv'_i, \bar{u}_iz_i, y_iv_i, y_iv'_i, y_iz_i, w_iv_i, w_iv'_i, w_iz_i, x_iv_i, x_iv'_i\}$. For each $j = 1, 2, \ldots, m$, corresponding to the clause $C_j = \{p_j, q_j, r_j\} \in \mathscr{C}$, associate a single vertex c_j and add edge set $E_j = \{c_jp_j, c_jq_j, c_jr_j\}$, $1 \le j \le m$. Finally, add a path $P = s_1s_2s_3$, join s_1 and s_3 to each vertex c_j with $1 \le j \le m$ and set $k = 1$.

Figure 1 shows an example of the graph obtained when $U = \{u_1, u_2, u_3, u_4\}$ and $\mathscr{C} = \{C_1, C_2, C_3\}$, where $C_1 = \{u_1, u_2, \bar{u}_3\}, C_2 = \{\bar{u}_1, u_2, u_4\}, C_3 = \{\bar{u}_2, u_3, u_4\}$.



Figure 1: An instance of the Roman bondage number problem resulting from an instance of 3SAT. Here $k = 1$ and $\gamma_{\rm R}(G) = 18$, where the bold vertex w means a Roman dominating function with $f(w) = 2$.

To prove that this is indeed a transformation, we only need to show that $b_{\rm R}(G)=1$ if and only if there is a truth assignment for U that satisfies all clauses in $\mathscr{C}$. This aim can be obtained by proving the following four claims.

Claim 4.1 $\gamma_{\rm R}(G)\geq 4n+2$. *Moreover, if* $\gamma_{\rm R}(G)=4n+2$, *then for any* $\gamma_{\rm R}$*-function* f *on* G, $f(H_i)=4$ *and at most one of* $f(u_i)$ *and* $f(\bar{u}_i)$ *is 2 for each* i, $f(c_j)=0$ *for each* j *and* $f(s_2)=2$.

Proof. Let f be a $\gamma_{\rm R}$-function of G, and let $H'_i=H_i-u_i-\bar{u}_i$.

If $f(u_i)=2$ and $f(\bar{u}_i)=2$, then $f(H_i)\geq 4$. Assume either $f(u_i)=2$ or $f(\bar{u}_i)=2$, if $f(x_i)=0$ or $f(y_i)=0$, then there is at least one vertex t in $\{v_i,\bar{v}_i,z_i\}$ such that $f(t)=2$. And hence $f(H'_i)\geq 2$. Thus, $f(H_i)\geq 4$.

If $f(u_i)\neq 2$ and $f(\bar{u}_i)\neq 2$, let f' be a restriction of f on H'_i, then f' is a Roman dominating function of H'_i, and $f'(H'_i)\geq\gamma_{\rm R}(H'_i)$. Since the maximum degree of H'_i is $V(H'_i)-3$, by Lemma 2.3, $\gamma_{\rm R}(H'_i)>3$ and hence $f'(H'_i)\geq 4$ and $f(H_i)\geq 4$. If $f(s_1)=0$ or $f(s_3)=0$, then there is at least one vertex t in $\{c_1,\cdots,c_m,s_2\}$ such that $f(t)=2$. Then $f(N_G[V(P)])\geq 2$, and hence $\gamma_{\rm R}(G)\geq 4n+2$.

Suppose that $\gamma_{\rm R}(G)=4n+2$, then $f(H_i)=4$ and since $f(N_G[x_i])\geq 1$, at most one of $f(u_i)$ and $f(\bar{u}_i)$ is 2 for each $i=1,2,\ldots,n$, while $f(N_G[V(P)])=2$. Then we have $f(s_2)=2$ since $f(N_G[s_2])\geq 1$. Consequently, $f(c_j)=0$ for each $j=1,2,\ldots,m$. ∎

Claim 4.2 $\gamma_{\rm R}(G)=4n+2$ *if and only if* $\mathscr{C}$ *is satisfiable.*

Proof. Suppose that $\gamma_{\rm R}(G)=4n+2$ and let f be a $\gamma_{\rm R}$-function of G. By Claim 4.1, at most one of $f(u_i)$ and $f(\bar{u}_i)$ is 2 for each $i=1,2,\ldots,n$. Define a mapping $t:U\to\{T,F\}$ by

$$t(u_i)=\begin{cases} T & \text{if } f(u_i)=2 \text{ or } f(u_i)\neq 2 \text{ and} f(\bar{u}_i)\neq 2,\\ F & \text{if } f(\bar{u}_i)=2.\end{cases}\qquad i=1,2,\ldots,n. \tag{4.1}$$

We now show that t is a satisfying truth assignment for $\mathscr{C}$. It is sufficient to show that every clause in $\mathscr{C}$ is satisfied by t. To this end, we arbitrarily choose a clause

$C_j \in \mathscr{C}$ with $1 \le j \le m$.

By Claim 4.1, $f(c_j) = f(s_1) = f(s_3) = 0$. There exists some i with $1 \le i \le n$ such that $f(u_i) = 2$ or $f(\bar{u}_i) = 2$ where c_j is adjacent to u_i or $\bar{u}_i$. Suppose that c_j is adjacent to u_i where $f(u_i) = 2$. Since u_i is adjacent to c_j in G, the literal u_i is in the clause C_j by the construction of G. Since $f(u_i) = 2$, it follows that $t(u_i) = T$ by (4.1), which implies that the clause C_j is satisfied by t. Suppose that c_j is adjacent to $\bar{u}_i$ where $f(\bar{u}_i) = 2$. Since $\bar{u}_i$ is adjacent to c_j in G, the literal $\bar{u}_i$ is in the clause C_j. Since $f(\bar{u}_i) = 2$, it follows that $t(u_i) = F$ by (4.1). Thus, t assigns $\bar{u}_i$ the truth value T, that is, t satisfies the clause C_j. By the arbitrariness of j with $1 \le j \le m$, we show that t satisfies all the clauses in $\mathscr{C}$, that is, $\mathscr{C}$ is satisfiable.

Conversely, suppose that $\mathscr{C}$ is satisfiable, and let $t : U \to \{T, F\}$ be a satisfying truth assignment for $\mathscr{C}$. Create a function f on $V(G)$ as follows: if $t(u_i) = T$, then let $f(u_i) = f(v'_i) = 2$, and if $t(u_i) = F$, then let $f(\bar{u}_i) = f(v_i) = 2$. Let $f(s_2) = 2$. Clearly, $f(G) = 4n + 2$. Since t is a satisfying truth assignment for $\mathscr{C}$, for each $j = 1, 2, \ldots, m$, at least one of literals in C_j is true under the assignment t. It follows that the corresponding vertex c_j in G is adjacent to at least one vertex w with $f(w) = 2$ since c_j is adjacent to each literal in C_j by the construction of G. Thus f is a Roman dominating function of G, and so $\gamma_{\mathrm{R}}(G) \le f(G) = 4n + 2$. By Claim 4.1, $\gamma_{\mathrm{R}}(G) \ge 4n + 2$, and so $\gamma_{\mathrm{R}}(G) = 4n + 2$. ∎

Claim 4.3 *$\gamma_{\mathrm{R}}(G - e) \le 4n + 3$ for any $e \in E(G)$.*

Proof. For any edge $e \in E(G)$, it is sufficient to construct a Roman dominating function f with weight $4n + 3$ of G. We first assume $e \in E_G(s_1)$ or $e \in E_G(s_3)$ or $e \in E_G(c_j)$ for each $j = 1, 2, \ldots, m$, without loss of generality let $e \in E_G(s_1)$ or $e = c_j u_i$ or $e = c_j \bar{u}_i$. Let $f(s_3) = 2, f(s_1) = 1$ and $f(u_i) = f(v'_i) = 2$ for each $i = 1, 2, \ldots, n$. For the edge $e \notin E_G(u_i)$ and $e \notin E_G(v'_i)$ or $e = \bar{u}_i z_i$, let $f(s_1) = 2, f(s_3) = 1$ and $f(u_i) = f(v'_i) = 2$. For the edge $e \notin E(\bar{u}_i)$ and $e \notin E(v_i)$ or $e = u_i z_i$, let $f(s_1) = 2, f(s_3) = 1$ and $f(\bar{u}_i) = f(v_i) = 2$. If $e = u_i v_i$ or $e = \bar{u}_i v'_i$, let $f(s_1) = 2, f(s_3) = 1$ and $f(x_i) = f(z_i) = 2$. Then f is a Roman dominating function of $G - e$ with $f(G - e) = 4n + 3$ and hence $\gamma_{\rm R}(G - e) \leq 4n + 3$. ∎

Claim 4.4 $\gamma_{\rm R}(G) = 4n + 2$ *if and only if* $b_{\rm R}(G) = 1$.

Proof. Assume $\gamma_{\rm R}(G) = 4n+2$ and consider the edge $e = s_1s_2$. Suppose $\gamma_{\rm R}(G) = \gamma_{\rm R}(G-e)$. Let f' be a $\gamma_{\rm R}$-function of $G-e$. It is clear that f' is also a $\gamma_{\rm R}$-function on G. By Claim 4.1 we have $f'(c_j) = 0$ for each $j = 1, 2, \ldots, m$ and $f'(s_2) = 2$. But then $f'(N_{G-e}[s_1]) = 0$, a contradiction. Hence, $\gamma_{\rm R}(G) < \gamma_{\rm R}(G - e)$, and so $b_{\rm R}(G) = 1$.

Now, assume $b_{\rm R}(G) = 1$. By Claim 4.1, we have that $\gamma_{\rm R}(G) \geq 4n+2$. Let e' be an edge such that $\gamma_{\rm R}(G) < \gamma_{\rm R}(G-e')$. By Claim 4.3, we have that $\gamma_{\rm R}(G-e') \leq 4n+3$. Thus, $4n + 2 \leq \gamma_{\rm R}(G) < \gamma_{\rm R}(G - e') \leq 4n + 3$, which yields $\gamma_{\rm R}(G) = 4n + 2$. ∎

By Claim 4.2 and Claim 4.4, we prove that $b_{\rm R}(G) = 1$ if and only if there is a truth assignment for U that satisfies all clauses in $\mathscr{C}$. Since the construction of the Roman bondage number instance is straightforward from a 3-satisfiability instance, the size of the Roman bondage number instance is bounded above by a polynomial function of the size of 3-satisfiability instance. It follows that this is a polynomial reduction.

The theorem follows. ∎

Corollary 4.1 *Roman domination number problem is NP-complete even for bipartite graphs.*

Proof. It is easy to see that the Roman bondage problem is in NP since a nondeterministic algorithm need only guess a vertex set pair (V_1, V_2) with $|V_1| + 2|V_2| \leq k$ and check in polynomial time whether that for any vertex $u \in V \setminus (V_1 \cup V_2)$ whether there is a vertex in V_2 adjacent to u for a given nonempty graph G.

We use the same method as Theorem 4.2 to prove this conclusion. We construct the same graph G but does not contain the path P. We set $k = 4n$, then use the same methods as Claim 4.1 and 4.2, we have that $\gamma_{\rm R}(G) = 4n$ if and only if $\mathscr{C}$ is satisfiable. ∎

5 General bounds

Lemma 5.1 *Let H be a spanning subgraph obtained by removing k edges from a graph G. Then $b_{\rm R}(G) \leq b_{\rm R}(H) + k$.*

Proof. Let $B = E(G) \setminus E(H)$ and B' be a minimum Roman bondage set of H. Then $|B| = k$, $|B'| = b_{\rm R}(H)$ and $\gamma_{\rm R}(H - B') > \gamma_{\rm R}(H)$. Let $f : V \to \{0, 1, 2\}$ be a Roman dominating function on H with $f(H) = \gamma_{\rm R}(H)$. Then each vertex x with $f(x) = 0$ is adjacent to at least one vertex y with $f(y) = 2$ in H, and so is in G since $H = G - B$, which implies that f is a Roman dominating function of G, and so $f(G) \geq \gamma_{\rm R}(G)$. It follows that $\gamma_{\rm R}(G - B - B') = \gamma_{\rm R}(H - B') > \gamma_{\rm R}(H) \geq \gamma_{\rm R}(G)$ and, hence, $b_{\rm R}(G) \leq |B| + |B'| = b_{\rm R}(H) + k$. ∎

Theorem 5.1 *$b_{\rm R}(G) \leq d_G(x) + d_G(y) + d_G(z) - |N_G(y) \cap N_G(\{x, z\})| - 3$ for any path (x, y, z) of length 2 in a graph G.*

Proof. Let $F_y = \{(y, u) \in E(G) : u \in N_G(y) \cap N_G(\{x, z\})\}$, $B = E_G(x) \cup E_G(z) \cup (E_G(y) \setminus F_y)$. Then

$$|B| = d_G(x) + d_G(y) + d_G(z) - |N_G(y) \cap N_G(\{x, z\})| - 2.$$

Let $H = G - B + yz$. Then x is an isolated vertex and z is a vertex of degree 1 which is only adjacent to y in H. Let f be a minimum Roman dominating function of H, then $f(x) = 1$ and $1 \leq f(y) + f(z) \leq 2$.

If $f(y) + f(z) = 2$, then let $f' = f$ except $f'(x) = 0$, $f'(y) = 2$ and $f'(z) = 0$. Clearly, f' is a Roman dominating function of G with $f'(G) < f(H)$ and, hence, $b_{\rm R}(G) \leq |B| - 1$.

If $f(y) + f(z) = 1$, then $f(y) = 0$ and $f(z) = 1$. There is an edge $(u, y) \in F_y$ with $f(u) = 2$. Let $f' = f$ except $f'(x) = 0$ if $u \in N_G(x)$ or $f'(z) = 0$ if $u \in N_G(z) \setminus N_G(x)$. Then f' is a Roman dominating function of G with $f'(G) < f(H)$, and hence $b_{\rm R}(G) \leq |B| - 1$. ∎

Theorem 5.2 *$b_{\rm R}(G) \leq d_G(x) + d_G(y) + d_G(z) - |N_G(y) \cap N_G(\{x, z\})| - |N_G(x) \cap N_G(z)| - 1$ for any path (x, y, z) of length 2 in a graph G.*

Proof. Let $F_y = \{(y,u) \in E(G) : u \in N_G(y) \cap N_G(\{x,z\})\}$ and $F_z = \{(z,u) \in E(G) : u \in (N_G(z) \cap N_G(x))\}$, $B = E_G(x) \cup (E_G(z) \setminus F_z) \cup (E_G(y) \setminus F_y)$ and $H = G - B$. Then x is an isolated vertex in H. Let f be a minimum Roman dominating function of H, then $f(x) = 1$. We will construct a Roman dominating function f' of G with $f'(G) < f(H)$.

If $f(z) = 0$, then there is an edge $(z,s) \in F_z$ with $f(s) = 2$. Thus, if $f(y) = 2$ or $f(z) = 0$, let $f' = f$ except $f'(x) = 0$. In the following, let $f(y) \neq 2$ and $f(z) \neq 0$.

If $f(y) = 0$. Then there is a vertex $s \in F_y$ such that $f(s) = 2$. If $s \in N_G(x)$, let $f' = f$ except $f'(x) = 0$. If $s \in N_G(z) \setminus N_G(x)$, let $f' = f$ except $f'(z) = 0$.

If $f(y) = 1$. If $f(z) = 1$, let $f' = f$ except $f'(x) = f'(z) = 0$ and $f'(y) = 2$. If $f(z) = 2$, let $f' = f$ except $f'(y) = 0$.

Then f' is a Roman dominating function of G with $f'(G) < f(H)$, and hence $b_{\rm R}(G) \leq |B| \leq d_G(x) + d_G(y) + d_G(z) - |N_G(y) \cap N_G(\{x,z\})| - |N_G(x) \cap N_G(z)| - 1$. ∎

Corollary 5.1 *$b_{\rm R}(G) \leq \min\{d_G(x) + d_G(y) + d_G(z) - |N_G(y) \cap N_G(\{x,z\})| - 3, d_G(x) + d_G(y) + d_G(z) - |N_G(y) \cap N_G(\{x,z\})| - |N_G(x) \cap N_G(z)| - 1\}$ for any path (x,y,z) of length 2 in a graph G.*

Corollary 5.2 *$b_{\rm R}(G) \leq 2\Delta(G) + \delta(G) - 3$ for any graph with diameter at least two.*

Corollary 5.3 *For any tree T of order at least 3, then $b_{\rm R}(T) \leq \Delta(T)$.*

Proof. If there is a vertex x adjacent to at least two vertices of degree one in T, say u_1 and u_2, then (u_1, x, u_2) is a path of length 2 in T. By Lemma 5.1, $b_{\rm R}(T) \leq d_T(u_1) + d_T(x) + d_T(u_2) - 3 \leq \Delta(T) - 1$.

Assume now that each vertex of T is adjacent to at most one vertex of degree one. Then T has a vertex u of degree 2 adjacent to exactly one vertex, say v, of degree one. Let w be the other vertex adjacent to u. Then (v,u,w) is a path of length 2 in T. By Lemma 5.1, $b_{\rm R}(T) \leq d_T(v) + d_T(u) + d_T(w) - 3 \leq \Delta(T)$. ∎

Lemma 5.2 *Let G be a connected graph of order n (≥ 3) and $\gamma_{\rm R}(G) = \gamma(G) + 1$. If there is an set B of edges with $\gamma_{\rm R}(G - B) = \gamma_{\rm R}(G)$, then $\Delta(G) = \Delta(G - B)$.*

Proof. Since G is connected and $n \geq 3$, $\gamma_{\rm R}(G) = \gamma(G) + 1 \leq n - 1$. Since $\gamma_{\rm R}(G - B) = \gamma_{\rm R}(G) \leq n - 1$, $G - B$ is nonempty. By Lemma 2.5 and Lemma 2.6, $\gamma_{\rm R}(G - B) \geq \gamma(G - B) + 1$. Since $\gamma_{\rm R}(G - B) = \gamma_{\rm R}(G) = \gamma(G) + 1 \leq \gamma(G - B) + 1$, $\gamma_{\rm R}(G - B) = \gamma(G - B) + 1$ and $\gamma(G - B) = \gamma(G)$.

If $G - B$ is connected, then by Lemma 2.7, $\Delta(G - B) = n - \gamma(G - B) = n - \gamma(G) = \Delta(G)$.

If $G - B$ is disconnected, then let G_1 be a nonempty connected component of $G - B$. By Lemma 2.5 and 2.6, $\gamma_{\rm R}(G_1) \geq \gamma(G_1) + 1$. Then $\gamma(G) + 1 = \gamma_{\rm R}(G - B) \geq \gamma_{\rm R}(G_1) + \gamma_{\rm R}(G - G_1) \geq \gamma(G_1) + 1 + \gamma(G - G_1) \geq \gamma(G) + 1$, thus $\gamma_{\rm R}(G_1) = \gamma(G_1) + 1$, $\gamma_{\rm R}(G - G_1) = \gamma(G - G_1)$ and $\gamma(G) = \gamma(G_1) + \gamma(G - G_1)$. By Lemma 2.6, $G - G_1$ is empty and hence $\gamma(G - G_1) = |V(G - G_1)|$. By Lemma 2.7, $\Delta(G_1) = |V(G_1)| - \gamma(G_1) = n - |V(G - G_1)| - \gamma(G_1) = n - \gamma(G - G_1) - \gamma(G_1) = n - \gamma(G) = \Delta(G)$. ∎

Theorem 5.3 *Let G be a connected graph of order n (≥ 3) and $\gamma_{\rm R}(G) = \gamma(G) + 1$. Then $b_{\rm R}(G) \leq \min\{b(G), n_\Delta\}$, where n_Δ is the number of vertices with maximum degree Δ in G.*

Proof. Since $n \geq 3$ and G is connected, $\Delta(G) \geq 2$ and hence $\gamma(G) \leq n - 2$. Let B be a minimum bondage set of G. Then $G - B$ is nonempty and by Lemma 2.5 and Lemma 2.6. Thus, $\gamma_{\rm R}(G - B) \geq \gamma(G - B) + 1 > \gamma(G) + 1 = \gamma_{\rm R}(G)$ and hence $b_{\rm R}(G) \leq b(G)$.

We now prove that $b_{\rm R}(G) \leq n_\Delta$. By Lemma 2.7, $\gamma_{\rm R}(G) = \gamma(G) + 1$ if and only if there is a vertex of degree $n - \gamma(G)$. If there is a vertex s in G such that $d_G(s) > n - \gamma(G)$, let $f(s) = 2$ and $f(w) = 1$ for any vertex w not in $N_G[s]$, then f is a Roman dominating function of G with $f(G) = \gamma(G)$, a contradiction. Thus, $\Delta(G) = n - \gamma(G)$. We can remove a smallest edge set B with $|B| \leq n_\Delta$ edges from G such that $\Delta(G - B) < \Delta(G) = n - \gamma(G)$ and $G - B$ is nonempty. Since $G - B$ is nonempty, by Lemma 2.5 and Lemma 2.6, $\gamma_{\rm R}(G - B) \geq \gamma(G - B) + 1$. Assume $\gamma_{\rm R}(G - B) = \gamma_{\rm R}(G)$, then by Lemma 5.2, $\Delta(G - B) = \Delta(G) = n - \gamma(G)$, a contradiction. Hence $b_{\rm R}(G) \leq |B| \leq n_\Delta$. ∎

Theorem 5.4 *For Roman graph G, $b_{\rm R}(G) \geq b(G)$.*

Proof. Let B be a minimum Roman bondage set of G, then $\gamma_{\rm R}(G-B) > \gamma_{\rm R}(G) = 2\gamma(G)$. By Lemma 2.5, $\gamma_{\rm R}(G-B) \le 2\gamma(G-B)$, then $\gamma(G-B) > \gamma(G)$ and hence $b_{\rm R}(G) \ge b(G)$. ∎

The equality in Theorem 5.4 can hold, for example, $b(C_{3k}) = 2 = b_{\rm R}(C_{3k})$, and the strict inequality can also hold, for example, $b(C_{3k+2}) = 2 < 3 = b_{\rm R}(C_{3k+2})$.

Theorem 5.5 *Let G be a nonempty graph with $\gamma_{\rm R}(G) \ge 3$. Then $b_{\rm R}(G) \le (\gamma_{\rm R}(G) - 2)\Delta(G) + 1$.*

Proof. The proof proceeds by induction on $\gamma_{\rm R}(G)$.

We first assume that $\gamma_{\rm R}(G) = 3$. Then by Lemma 2.3, $\Delta(G) = |V(G)| - 2$. Assume that $b_{\rm R}(G) \ge \Delta(G)+2$. Let u be a vertex of maximum degree in G. We have $\gamma_{\rm R}(G-u) = \gamma_{\rm R}(G) - 1 = 2$. There is a vertex v that is adjacent to every vertex in $G-u$ and hence $vu \notin E(G)$. Since $b_{\rm R}(G-u) \ge 2$, then for any edge $e \in E_{G-u}(v)$, $\gamma_{\rm R}(G-u-e) = 2$. Thus there is a vertex w that is adjacent to every vertex of $G-u-e$. But, since v is the only vertex of G that is not adjacent to u, $wu \in E(G)$, $d_G(w) = |V(G)| - 1$, a contradiction. Thus, $b_{\rm R}(G) \le \Delta(G) + 1$ if $\gamma_{\rm R}(G) = 3$.

Assume the induction hypothesis for any integer k and any graph H with $\gamma_{\rm R}(H) = k \ge 3$. Let G be a nonempty graph with $\gamma_{\rm R}(G) = k+1$, and assume that $b_{\rm R}(G) \ge (k-1)\Delta(G) + 2$. For any vertex u of G, let $H = G-u$. Then, $\gamma_{\rm R}(H) = \gamma_{\rm R}(G) - 1 = k$ since $d_G(u) < b_{\rm R}(G)$. By the inductive hypothesis and by Lemma 5.1, we have

$$\begin{aligned} b_{\rm R}(G) &\le b_{\rm R}(H) + d_G(u) \\ &\le (k-2)\Delta(H) + 1 + d_G(u) \\ &\le (k-2)\Delta(G) + 1 + \Delta(G) \\ &= (k-1)\Delta(G) + 1, \end{aligned}$$

a contradiction. Thus, $b_{\rm R}(G) \le (k-1)\Delta(G) + 1$, and by the principle of mathematical induction, $b_{\rm R}(G) \le (\gamma_{\rm R}(G) - 2)\Delta(G) + 1$. ∎

Use $\kappa(G)$ (resp. $\lambda(G)$) to denote the vertex-connectivity (resp. the edge-connectivity) of a connected graph G which is the minimum number of vertices (resp, edges) whose removal results in G disconnected. The famous Whitney's inequality can be stated as $\kappa(G) \leqslant \lambda(G) \leqslant \delta(G)$ for any graph G. A subset $F \subseteq E(G)$ is called a λ-cut if $|F| = \lambda(G)$ and $G-F$ is disconnected.

Theorem 5.6 *If G is a connected graph with order at least 3, then $b_{\rm R}(G) \leq 2\Delta(G) + \lambda(G) - 3$, where $\lambda(G)$ is the edge-connectivity of G.*

Proof. Let G be a connected graph with edge-connectivity $\lambda(G)$ and F be λ-cut of G. Then $H = G - F$ has exact two connected components. Let $x, y \in V(G)$, $xy \in F$, and H_x and H_y denote the components of $G - F$ containing x and y, respectively. Without loss of generality, let z be adjacent to x in H_x since $|V(G)| \geq 3$. Let $B = F \cup E_{H_x}(x) \cup E_{H_x}(z) - xz$ and f be a $\gamma_{\rm R}$-function of $G' = G - B$. Then x and z is only adjacent to each other in G', and so we can assume $f(x) = 2$ and $f(z) = 0$. We construct a Roman dominating function f' of G with $f'(G) < f(G')$.

If $V(H_y) = \{y\}$, then $f(y) = 1$. Let $f' = f$ except $f'(y) = 0$. Then f' is a Roman dominating function of G with $f'(G) < f(G')$. Thus, $b_{\rm R}(G) \leq |B| \leq 2\Delta(G) + \lambda(G) - 3$. In the following, we assume $|V(H_y)| \geq 2$.

If $\gamma_{\rm R}(H_y - y) \geq \gamma_{\rm R}(H_y)$, then

$$\gamma_{\rm R}(G - (F \cup E_{H_y}(y))) \geq \gamma_{\rm R}(H_x) + \gamma_{\rm R}(H_y) + 1 \geq \gamma_{\rm R}(G) + 1.$$

Thus

$$\begin{aligned} b_{\rm R}(G) &\leq |F \cup E_{H_y}(y))| \leq \Delta(G) + \lambda(G) - 1 \\ &\leq 2\Delta(G) + \lambda(G) - 3. \end{aligned}$$

If $\gamma_{\rm R}(H_y - y) = \gamma_{\rm R}(H_y) - 1$, we can assume that $f(y) = 1$. Let $f' = f$ except $f'(y) = 0$. Then f' is a Roman dominating function of G with $f'(G) < f(G')$. Thus,

$$b_{\rm R}(G) \leq |B| \leq 2\Delta(G) + \lambda(G) - 3.$$

The theorem follows. ∎

Considering vertex rather than edge-connectivity, we could conjecture an analogy of Theorem 5.6 by a similar argument.

Conjecture 5.1 *If G is a connected graph with order no less than 3, then $b_{\rm R}(G) \leq 2\Delta(G) + \kappa(G) - 3$, where $\kappa(G)$ is the vertex-connectivity of G.*

Theorem 5.7 *If G is a nonempty graph with a unique minimum Roman dominating function, then $b_{\rm R}(G) = 1$.*

Proof. Let f be the unique $\gamma_{\rm R}$-function on G, and let x be a vertex in G with $f(x) = 0$. Then there is a vertex $y \in N_G(x)$ with $f(y) = 2$. If there are at least two vertices $y, z \in N_G(x)$ such that $f(y) = f(z) = 2$ for each vertex x with $f(x) = 0$. Then let $f' = f$ except that $f'(x) = 2$ and $f'(y) = 0$ and f' is a $\gamma_{\rm R}$-function on G as well, which is a contradiction to the uniqueness of f. Thus, there is a unique $y \in N_G(x)$ with $f(y) = 2$ for a vertex x with $f(x) = 0$. Then $\gamma_{\rm R}(G - xy) > \gamma_{\rm R}(G)$, which implies that $b_{\rm R}(G) = 1$. ▮

Theorem 5.8 *If G is a vrc-graph with $\gamma_{\rm R}(G) = 3$, then $b_{\rm R}(G) \leq \Delta(G) + 1$.*

Proof. By Lemma 2.9, G is a vc-graph with $\gamma(G) = 2$. By Lemma 2.8, G is a complete $K_{2t}(t \geq 2)$ with a perfect matching M removed. Thus, G is $\Delta(G)$-regular, where $\Delta(G) = 2t - 2$. Let $uv \in M$. Then v is the only vertex not adjacent to u in G. Let $H = G - u$. Then $\gamma_{\rm R}(H) = 2$ since G is a vrc-graph with $\gamma_{\rm R}(G) = 3$. Note that the vertex v is the only vertex adjacent to all the other vertices in H adjacent to each of other vertices in H. Thus H has a unique minimum Roman dominating function f with $f(v) = 2 = \gamma_{\rm R}(H)$. By Theorem 5.7, $b_{\rm R}(H) = 1$ and hence $b_{\rm R}(G) \leq \Delta(G) + 1$. ▮

Theorem 5.9 *If there exists at least one vertex u in a graph G with $\gamma_{\rm R}(G-u) \geq \gamma_{\rm R}(G)$, then $b_{\rm R}(G) = d_G(x) \leq \Delta(G)$.*

Proof. Since $\gamma_{\rm R}(G - E_G(u)) = \gamma_{\rm R}(G - u) + 1 > \gamma_{\rm R}(G)$, $b_{\rm R}(G) = d_G(x) \leq \Delta(G)$. ▮

Corollary 5.4 *Let G be a graph of order n. If $\gamma_{\rm R}(G) = 3 \neq n$, then $b_{\rm R}(G) \leq \Delta + 1$.*

Problem 5.1 *Whether or not there exits a positive integer c such that $b_{\rm R}(G) \leq \Delta(G)+c$ for any graph G of order n and $\gamma_{\rm R}(G) \neq n$.*

The *vertex covering number* $\beta(G)$ of G is the minimum number of vertices that are incident with all edges in G. If G has no isolated vertices, then $\gamma_{\rm R}(G) \leq 2\gamma(G) \leq 2\beta(G)$. If $\gamma_{\rm R}(G) = 2\beta(G)$, then $\gamma_{\rm R}(G) = 2\gamma(G)$ and hence G is a Roman graph. In [17], Volkmann gave a lot of graphs with $\gamma(G) = \beta(G)$.

Theorem 5.10 *Let G be a graph with $\gamma_{\mathrm{R}}(G) = 2\beta(G)$. Then*

(1) $b_{\mathrm{R}}(G) \geq \delta(G)$*;*

(2) $b_{\mathrm{R}}(G) \geq \delta(G) + 1$ *if G is a vrc-graph.*

Proof. Let G be a graph with $\gamma_{\mathrm{R}}(G) = 2\beta(G)$.

(1) Without loss of generality, Assume $\delta(G) \geq 2$. Let $B \subseteq E(G)$ with $|B| \leq \delta(G) - 1$. Then $\delta(G - B) \geq 1$ and so $\gamma_{\mathrm{R}}(G) \leq \gamma_{\mathrm{R}}(G - B) \leq 2\beta(G - B) \leq 2\beta(G) = \gamma_{\mathrm{R}}(G)$. Thus, B is not a Roman bondage set of G, and so $b_{\mathrm{R}}(G) \geq \delta(G)$.

(2) From the above proof, every Roman bondage set B contains at least all edges incident with some vertex x, so that $G - B$ has an isolated vertex. On the other hand, if G is a vrc-graph, then $\gamma_{\mathrm{R}}(G - x) < \gamma_{\mathrm{R}}(G)$ for any vertex x, which implies that the removal of all edges incident with x can not enlarge the Roman domination number. Hence $b_{\mathrm{R}}(G) \geq \delta(G) + 1$. ∎

References

[1] R. C. Brigham, P. Z. Chinn and R. D. Dutton, Vertex domination-cirtical graphs. *Networks*, **18** (1988), 173-179.

[2] E. J. Cockayne, P. A. Dreyer, Jr., S. M. Hedetniemi, S. T. Hedetniemi, A. A. McRae, Roman domination in graphs. *Discrete Mathematics*, **278** (1-3) (2004), 11-22.

[3] E. W. Chambers, B. Kinnersley, N. Prince, D. B. West, Extremal problems for Roman domination. *SIAM J. Discrete Math.* **23** (2009), 1575-1586.

[4] J. F. Fink, M. S. Jacobson, L. F. Kinch and J. Roberts, The bondage number of a graph. *Discrete Mathematics*, **86** (1990), 47-57.

[5] O. Favaron, H. Karami, R. Khoeilar and S. M. Sheikholeslami, On the Roman domination number of a graph. *Discrete Mathematics*, **309** (2009), 3447-3451.

[6] X. L. Fu, Y. S. Yang and B. Q. Jiang, Roman domination in regular graphs. *Discrete Mathematics*, **309** (2009), 1528-1537.

[7] M. R. Garey, D. S. Johnson, *Computers and Intractability: A Guide to the Theory of NP-Completeness*, Freeman, San Francisco, 1979.

[8] T. W. Haynes, S. T. Hedetniemi, P. J. Slater, *Fundamentals of Domination in Graphs*, Marcel Dekker, New York, 1997.

[9] T. W. Haynes, S. T. Hedetniemi, P. J. Slater, *Domination in Graphs: Advanced Topics*, Marcel Dekker, New York, 1997.

[10] J. H. Hattingh, A. R. Plummer, Restrained bondage in graphs. *Discrete Mathematics*, **308** (2008), 5446-5453.

[11] M. Liedloff, T. Kloks, J. P. Liu and S. L. Peng, Efficient algorithms for Roman domination on some classes of graphs, *Discrete Applied Mathematics*, **156** (2008), 3400-3415.

[12] M. Liedloff, T. Kloks, J. P. Liu and S. L. Peng, Roman domination over some graph classes, *Lecture Notes in Computer Science*, 2005, Volume **3787**, Graph-Theoretic Concepts in Computer Science, Pages 103-114.

[13] A. Pagourtzis, P. Penna, K. Schlude, K. Steinhofel, D. Taylor and P. Widmayer, Server placements, Roman domination and other dominating set variants, *in Proc. Second International Conference on Theoretical Computer Science* (2002), 280-291.

[14] R. R. Rubalcaba, P. J. Slater, Roman dominating influence parameters. *Discrete Mathematics*, **307** (2007), 3194-3200.

[15] W. P. Shang and X. D. Hu, The roman domination problem in unit disk graphs, *Lecture Notes in Computer Science*, 2007, Volume 4489/2007, 305-312.

[16] W. P. Shang and X. D. Hu, Roman domination and its variants in unit disk graphs, *Discrete Mathematics, Algorithms and Applications*, **2** (2010), 99-105.

[17] L. Volkmann, On graphs with equal domination and covering numbers. *Discrete Applied Mathematics*, **51** (1994), 211-217.

[18] J.-M. Xu, *Toplogical Structure and Analysis of Interconnection Networks*. Kluwer Academic Publishers, Dordrecht/Boston/London, 2001.

[19] J.-M. Xu, *Theory and Application of Graphs*. Kluwer Academic Publishers, Dordrecht/Boston/London, 2003.

[20] H. M. Xing, X. Chen and X. G. Chen, A note on Roman domination in graphs. *Discrete Mathematics*, **306** (2006), 3338-3340.